

**This Page Is Inserted by IFW Operations
and is not a part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- **BLACK BORDERS**
- **TEXT CUT OFF AT TOP, BOTTOM OR SIDES**
- **FADED TEXT**
- **ILLEGIBLE TEXT**
- **SKEWED/SLANTED IMAGES**
- **COLORED PHOTOS**
- **BLACK OR VERY BLACK AND WHITE DARK PHOTOS**
- **GRAY SCALE DOCUMENTS**

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵: C12N 9/76, 15/57, C11D 3/386	A1	(11) International Publication Number: WO 94/25583 (43) International Publication Date: 10 November 1994 (10.11.94)
(21) International Application Number: PCT/DK94/00177 (22) International Filing Date: 4 May 1994 (04.05.94) (30) Priority Data: 0523/93 5 May 1993 (05.05.93) DK (71) Applicant (for all designated States except US): NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsværd (DK). (72) Inventors; and (75) Inventors/Applicants (for US only): BRANNER, Sven [DK/DK]; Ved Smedebakken 7A, DK-2800 Lyngby (DK). HASTRUP, Sven [DK/DK]; Frederiksborgvej 10, 3.th., DK-2400 Copenhagen NV (DK). (74) Common Representative: NOVO NORDISK A/S; Att.: Lise Abildgaard Sørensen, Corporate Patents, Novo Allé, DK-2880 Bagsværd (DK).		(81) Designated States: AU, BB, BG, BR, BY, CA, CN, CZ, FI, HU, JP, KP, KR, KZ, LK, LV, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: A RECOMBINANT TRYPSIN-LIKE PROTEASE (57) Abstract An active recombinant trypsin-like protease enzyme comprising the amino acid residues 25-224 of the amino acid sequence shown in the appended SEQ ID No. 2 as well as a DNA construct encoding said enzyme. The enzyme may be used as a detergent enzyme.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LJ	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

A recombinant trypsin-like protease

FIELD OF THE INVENTION

The present invention relates to an active recombinant trypsin-like protease, a proteolytic composition comprising said trypsin-like protease, a DNA construct
5 comprising a DNA sequence encoding said trypsin-like protease, and a vector and cell harbouring the DNA construct. Furthermore, the present invention relates to a method of preparing the trypsin-like protease by use of recombinant DNA techniques.

BACKGROUND OF THE INVENTION

10 Proteases are widely used as ingredients in commercial detergents to improve the detergency towards proteinaceous soiling. For a number of years proteases with a broad specificity, such as subtilisins (*Bacillus* serine proteases) have been the most widely used detergent proteases. However, in recent years proteases having a more narrow specificity have been of
15 increasing interest for detergent purposes.

WO 89/06270 discloses a detergent composition comprising a protease with a narrow substrate specificity, namely a trypsin-like protease capable of cleaving peptide bonds at the C-terminal side of lysine or arginine. The trypsin-like protease disclosed in said reference is produced by conventional fermentation
20 of a strain of the *Fusarium* sp. *F. oxysporum* DSM 2672, and is inevitably produced together with another protease (Protease I), the existence of which is undesirable in detergent compositions. Thus, production of the trypsin-like protease from the *F. oxysporum* strain involves a step in which the undesired protease is separated from the trypsin-like protease, which makes the large
25 scale production of the latter protease more expensive than what is desirable.

It would be desirable to facilitate the production of said *Fusarium* trypsin-like protease, in particular by avoiding the co-production of the above mentioned undesired Protease I, to be able of producing both larger amounts of the enzyme and to produce it in a more economical manner than what is possible by the prior art methods.

BRIEF DISCLOSURE OF THE INVENTION

The present inventors have now succeeded in cloning a DNA sequence encoding a *Fusarium* trypsin-like protease and in obtaining expression of an active trypsin-like protease from said DNA sequence.

10 Accordingly, in a first aspect the present invention relates to an active recombinant trypsin-like protease comprising the amino acid residues 1-224 of the amino acid sequence shown in the appended SEQ ID No. 2.

In the present context the term "trypsin-like protease" is intended to indicate an enzyme having an activity similar to that of trypsin, i.e. an enzyme capable
15 of cleaving peptide bonds at the C-terminal side of lysine or arginine. The trypsin-like protease activity may be determined in an assay based on cleavage of a trypsin substrate such as N-Benzoyl-L-arginine-p-nitroanilide hydrochloride (L-BAPA or L-BAPNA), e.g. as described in the Materials and Methods section below.

20 The term "recombinant" as used about the trypsin-like protease of the invention is intended to indicate that it is produced by a cell transformed with a DNA sequence encoding the protease. Thus, the recombinant trypsin-like protease is produced by another organism than its parent organism and accordingly essentially free from components derived from said parent
25 organism, i.e. components produced by the *F. oxysporum* strain DSM 2672. Such components may confer undesirable properties to the protease, for in-

stance by giving rise to undesirable enzymatic activities. In this respect, the recombinant trypsin-like protease of the invention is easily produced without Protease I, and no costly separation similar to that performed according to WO 89/06270 is required.

5 In the course of the research leading to the present invention, which is described in detail in the following examples, it was quite unexpectedly found to be difficult to obtain any substantial production of an active trypsin-like protease by cultivation of a conventionally used production organism (*Aspergillus oryzae*) transformed with the DNA sequence shown in SEQ ID No.
10 1. On the basis of analyses of the DNA sequence and the expression product, it was found that the enzyme encoded by the DNA sequence was expressed as an inactive proenzyme, and that large amounts of this proenzyme were expressed. Thus, even though large amounts of the precursor form of the active enzyme were available no substantial yield of an active matured
15 enzyme could be obtained, and it was concluded that the proenzyme was unstable in the *A. oryzae* fermentation broth.

It was surprisingly found that the yield of the active recombinant trypsin-like protease could be substantially increased if the medium, in which a host cell harbouring a DNA sequence encoding a proenzyme form of the enzyme was
20 cultivated, also comprised a proteolytic enzyme capable of converting the proenzyme into an active enzyme in situ. It is expected that the low enzymatic activity expressed when no other proteolytic enzyme is present may be ascribed to the fact that the inactive proenzyme form of the trypsin-like protease is considerably unstable and degraded rapidly by proteolytic enzymes
25 expressed from the host organism, and that the addition of a proteolytic enzyme, which more readily than proenzyme-degrading proteolytic enzymes, reacts with the proenzyme resulting in the production of a mature, active trypsin-like protease.

In a further aspect the present invention relates to a DNA construct comprising a DNA sequence encoding an active recombinant trypsin-like protease of the invention as defined above.

In still further aspects the present invention relates to a recombinant
5 expression vector harbouring the DNA construct of the invention, to a cell which either harbours the DNA construct or the expression vector of the invention, and to a process for the production of a trypsin-like protease of the invention, wherein a cell of the invention as described above is cultured under conditions conducive to the production of the protease, and the protease is
10 subsequently recovered from the culture.

Finally, the present invention relates to a detergent additive and detergent composition comprising the active trypsin-like protease of the invention.

DETAILED DESCRIPTION OF THE INVENTION

The amino acid sequence of the recombinant trypsin-like protease of the
15 invention, which was isolated from a strain of the *Fusarium oxysporum* DSM 2672, has been aligned with that of trypsin-like proteases of other origins and have been shown to have a degree of homology of about 43% with that of bovine trypsin (GenBank Acc. No. P00760) and 44% with that of *Streptomyces griseus* (GenBank Acc. No. P00776). The highest homology of
20 the trypsin-like protease of the invention has been observed with that of the fruit fly (GenBank Acc. No. A23493), namely 48%.

These homologies are taken to indicate that some kind of evolutionary relationship exists between trypsin-like proteases, but also that the *Fusarium* trypsin-like protease of the invention may represent a distinct class of trypsin-
25 like proteases. It is contemplated that the trypsin-like protease of the invention or DNA encoding the protease may be isolated from other organisms, including

animals, especially a mammal, an insect, a plant or a microorganism. In the present context, especially interesting origins are bacteria and fungi, including yeasts and filamentous fungi.

The active recombinant trypsin-like protease of the invention comprising the 5 amino acid residues 1-224 of the SEQ ID No. 2 has a number of characteristic properties. For instance, the protease surprisingly shows a reversed Arg/Lys specificity compared to that of bovine trypsin, i.e. the trypsin-like protease of the invention is more Arg-active than Lys-active.

The pH-activity profile of the trypsin-like protease of the invention shows a 10 broad activity optimum between pH 8 and 11, in particular about 10 when using H-D-Val-Leu-Lys-pNA as a substrate at 25°C. The temperature optimum is approximately 40°C at pH 9.5 using H-D-Val-Leu-Lys-pNA. The mature, active enzyme consists of 224 amino acid residues and has a molecular weight of 22190. It is expressed in the form of a pre-pro-enzyme, in which 15 the signal peptide is contemplated to be amino acid residues -24 to -8 (according to the rules of von Heijne (1986) and the pro-peptide amino acid residues -7 to -1. The specific activity of the enzyme is at least about 30 Casein Protease Units (CPU)/g and preferably equal to or higher than about 35 CPU/g. The protease activity may be determined by the method given in WO 20 89/06270, the content of which is hereby incorporated by reference.

The DNA sequence of the DNA construct of the invention encoding a recombinant protease enzyme as defined above is preferably as shown in the appended SEQ ID No. 1. Analogues of said sequence, which differ in one or more codons, but which encodes the recombinant protease protein are also 25 within the invention.

The DNA sequence of the DNA construct of the invention may be isolated by well-known methods. Thus, the DNA sequence may, for instance, be isolated by establishing a cDNA or genomic library from an organism expected to

harbour the sequence, and screening for positive clones by conventional procedures. Examples of such procedures are hybridization to oligonucleotide probes synthesized on the basis of the full amino acid sequence shown in SEQ ID No. 2 or a subsequence thereof in accordance with standard techniques (cf. 5 Sambrook et al., 1989), and/or selection for clones expressing a trypsin-like protease activity as defined above, and/or selection for clones producing a protein which is reactive with an antibody raised against the trypsin-like protease comprising the amino acid sequence shown in SEQ ID No. 2 and in particular amino acid residues 1-224 thereof.

10 A preferred method of isolating a DNA construct of the invention from a cDNA or genomic library is by use of polymerase chain reaction (PCR) using degenerate oligonucleotide probes prepared on the basis of the amino acid sequence of the trypsin-like protease of the invention comprising amino acid residues 1-224 of SEQ ID No. 2. For instance, the PCR may be carried out 15 using the techniques described in US Patent No. 4,683,202 or by R.K. Saiki et al. (1988).

Alternatively, the DNA sequence of the DNA construct of the invention may be prepared synthetically by established standard methods, e.g. the phosphoramidite method described by Beaucage and Caruthers (1981), or the 20 method described by Matthes et al. (1984). According to the phosphoramidite method, oligonucleotides are synthesized, e.g. in an automatic DNA synthesizer, purified, annealed, ligated and cloned in appropriate vectors.

Finally, the DNA construct may be of mixed genomic and synthetic, mixed synthetic and cDNA or mixed genomic and cDNA origin prepared by ligating 25 fragments of synthetic, genomic or cDNA origin (as appropriate), the fragments corresponding to various parts of the entire recombinant DNA molecule, in accordance with standard techniques.

The recombinant expression vector carrying the DNA construct of the invention may be any vector which may conveniently be subjected to recombinant DNA procedures, and the choice of vector will often depend on the host cell into which it is to be introduced. Thus, the vector may be an
5 autonomously replicating vector, i.e. a vector which exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g. a plasmid, a bacteriophage or an extrachromosomal element, minichromosome or an artificial chromosome. Alternatively, the vector may be one which, when introduced into a host cell, is integrated into the host cell
10 genome and replicated together with the chromosome(s) into which it has been integrated.

In the vector, the DNA sequence should be operably connected to a suitable promoter sequence. The promoter may be any DNA sequence which shows transcriptional activity in the host cell of choice and may be derived from
15 genes encoding proteins either homologous or heterologous to the host cell. Examples of suitable promoters for directing the transcription of the DNA construct of the invention, especially in a bacterial host, are the promoter of the *lac* operon of *E.coli*, the *Streptomyces coelicolor* agarase gene *dagA* promoters, the promoters of the *Bacillus licheniformis* α -amylase gene (*amyL*),
20 the promoters of the *Bacillus stearothermophilus* maltogenic amylase gene (*amyM*), the promoters of the *Bacillus Amyloliquefaciens* α -amylase (*amyQ*), the promoters of the *Bacillus subtilis* *xylA* and *xylB* genes etc. For transcription in a fungal host, examples of useful promoters are those derived from the gene encoding *A. oryzae* TKA amylase, *Rhizomucor miehei* aspartic
25 proteinase, *A. niger* neutral α -amylase, *A. niger* acid stable α -amylase, *A. niger* glucoamylase, *Rhizomucor miehei* lipase, *A. oryzae* alkaline protease, *A. oryzae* triose phosphate isomerase or *A. nidulans* acetamidase.

The expression vector of the invention may also comprise a suitable transcription terminator and, in eukaryotes, polyadenylation sequences
30 operably connected to the DNA sequence encoding the recombinant protease

of the invention. Termination and polyadenylation sequences may suitably be derived from the same sources as the promoter.

The vector may further comprise a DNA sequence enabling the vector to replicate in the host cell in question. Examples of such sequences are the origins of replication of plasmids pUC19, pACYC177, pUB110, pE194, pAMB1 and pIJ702.

The vector may also comprise a selectable marker, e.g. a gene the product of which complements a defect in the host cell, such as the *dal* genes from *B.subtilis* or *B.licheniformis*, or one which confers antibiotic resistance such as ampicillin, kanamycin, chloramphenicol or tetracyclin resistance. Examples of *Aspergillus* selection markers include amdS, argB, niaD and sC, a marker giving rise to hygromycin resistance. Furthermore, the selection may be accomplished by co-transformation, e.g. as described in WO 91/17243.

While intracellular expression may be advantageous in some respects, e.g. when using certain bacteria as host cells, it is generally preferred that the expression is extracellular. As mentioned above the trypsin-like protease of the invention comprising the amino acid sequence shown in the SEQ ID No. 2 comprises a preregion permitting secretion of the expressed protease into the culture medium. If desirable, this preregion may be substituted with a different preregion or signal sequence, convenient accomplished by substitution of the DNA sequences encoding the respective preregions.

The procedures used to ligate the DNA construct of the invention, the promoter, terminator and other elements, respectively, and to insert them into suitable vectors containing the information necessary for replication, are well known to persons skilled in the art (cf., for instance, Sambrook et al. (1989)).

The cell of the invention either comprising a DNA construct or an expression vector of the invention as defined above is advantageously used as a host cell

in the recombinant production of a polypeptide of the invention. The cell may be transformed with the DNA construct of the invention, conveniently by integrating the DNA construct in the host chromosome. This integration is generally considered to be an advantage as the DNA sequence is more likely
5 to be stably maintained in the cell. Integration of the DNA constructs into the host chromosome may be performed according to conventional methods, e.g. by homologous or heterologous recombination. Alternatively, the cell may be transformed with an expression vector as described above in connection with the different types of host cells.

- 10 The cell of the invention may be a cell of a higher organism such as a mammal or an insect, but is preferably a microbial cell, e.g. a bacterial or a fungal (including yeast) cell.

Examples of suitable bacteria are grampositive bacteria such as *Bacillus subtilis*, *Bacillus licheniformis*, *Bacillus lentus*, *Bacillus brevis*, *Bacillus*
15 *stearothermophilus*, *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus coagulans*, *Bacillus circulans*, *Bacillus lautus*, *Bacillus megaterium*, *Bacillus thuringiensis*, or *Streptomyces lividans* or *Streptomyces murinus*, or gramnegative bacteria such as *E.coli*. The transformation of the bacteria may for instance be effected by protoplast transformation or by using competent cells
20 in a manner known *per se*.

The yeast organism may favourably be selected from a species of *Saccharomyces* or *Schizosaccharomyces*, e.g. *Saccharomyces cerevisiae*. The filamentous fungus may advantageously belong to a species of *Aspergillus*, e.g. *Aspergillus oryzae* or *Aspergillus niger*. Alternatively, a strain of a
25 *Fusarium* species, e.g. *F. oxysporum*, can be used as a host cell. Fungal cells may be transformed by a process involving protoplast formation and transformation of the protoplasts followed by regeneration of the cell wall in a manner known *per se*. A suitable procedure for transformation of *Aspergillus*

host cells is described in EP 238 023. A suitable method of transforming *Fusarium* species is described by Malardier et al., 1989.

In a yet further aspect, the present invention relates to a method of producing a recombinant trypsin-like protease of the invention, which method comprises
5 cultivating a host cell as described above under conditions conducive to the production of the protease and recovering the protease from the cells and/or culture medium.

The medium used to cultivate the cells may be any conventional medium suitable for growing the host cell in question and obtaining expression of the
10 protease of the invention. Suitable media are available from commercial suppliers or may be prepared according to published recipes (e.g. in catalogues of the American Type Culture Collection).

From the above disclosure it is apparent that when the trypsin-like protease of the invention is expressed as a proenzyme, substantially increased yields may
15 be obtained when the cultivation is performed in the presence of a proteolytic enzyme or optionally two or more proteolytic enzymes capable of transforming the proenzyme into an active enzyme in the medium.

The proteolytic enzyme(s) may be added as such to the fermentation broth in which the cell producing the proenzyme is cultured. The proteolytic enzyme(s)
20 to be added may be a *Bacillus* metallo-protease, for instance a commercially available enzyme, e.g. Neutrase®, thermolysin, or another enzyme with a similar activity such as a proteolytic enzyme from *Bacillus stearothermophilus* (Zamost et al., 1990) or an aspartic protease from *F. oxysporum* obtainable as described in Example 1 hereinafter. Alternatively, the proteolytic enzyme may
25 be produced by culturing a host cell transformed with a DNA sequence encoding the proteolytic enzyme under suitable conditions to produce the enzyme and recovering the enzyme from the culture.

Furthermore, the presence of the proteolytic enzyme(s) in the fermentation broth may be accomplished by inserting the DNA sequence(s) encoding said enzyme(s) into the cell harbouring a DNA construct of the invention in such a manner that the proteolytic enzyme(s) is/are expressed and excreted in sufficient amounts to activate the protease of the invention.

The resulting trypsin-like protease may be recovered from the medium by conventional procedures including separating the cells from the medium by centrifugation or filtration, if necessary after disruption of the cells, precipitating the proteinaceous components of the supernatant or filtrate by means of a salt, e.g. ammonium sulphate, followed by purification by a variety of chromatographic procedures, e.g. ion exchange chromatography, affinity chromatography, or the like.

Detergent Compositions

According to the invention, the trypsin-like protease may typically be a component of a detergent composition. As such, it may be included in the detergent composition in the form of a non-dusting granulate, a stabilized liquid, or a protected enzyme. Non-dusting granulates may be produced, e.g., as disclosed in US 4,106,991 and 4,661,452 (both to Novo Industri A/S) and may optionally be coated by methods known in the art. Examples of waxy coating materials are poly(ethylene oxide) products (polyethyleneglycol, PEG) with mean molar weights of 1000 to 20000, ethoxylated nonylphenols having from 16 to 50 ethylene oxide units; ethoxylated fatty alcohols in which the alcohol contains from 12 to 20 carbon atoms and in which there are 15 to 80 ethylene oxide units; fatty alcohols; fatty acids; and mono- and di- and triglycerides of fatty acids. Examples of film-forming coating materials suitable for application by fluid bed techniques are given in patent GB 1483591. Liquid enzyme preparations may, for instance, be stabilized by adding a polyol such as propylene glycol, a sugar or sugar alcohol, lactic acid or boric acid according to established methods. Other enzyme stabilizers are well known in

the art. Protected enzymes may be prepared according to the method disclosed in EP 238,216.

The detergent composition of the invention may be in any convenient form, e.g. as powder, granules, paste or liquid. A liquid detergent may be aqueous, 5 typically containing up to 70 % water and 0-30 % organic solvent, or nonaqueous.

The detergent composition comprises one or more surfactants, each of which may be anionic, nonionic, cationic, or zwitterionic. The detergent will usually contain 0-50 % of anionic surfactant such as linear alkylbenzenesulfonate 10 (LAS), alpha-olefinsulfonate (AOS), alkyl sulfate (fatty alcohol sulfate) (AS), alcohol ethoxysulfate (AEOS or AES), secondary alkanesulfonates (SAS), alpha-sulfo fatty acid methyl esters, alkyl- or alkenylsuccinic acid or soap. It may also contain 0-40 % of nonionic surfactant such as alcohol ethoxylate (AEO or AE), carboxylated alcohol ethoxylates, nonylphenol ethoxylate, 15 alkylpolyglycoside, alkyltrimethylamineoxide, ethoxylated fatty acid monoethanolamide, fatty acid monoethanolamide, or polyhydroxy alkyl fatty acid amide (e.g. as described in WO 92/06154).

The detergent composition may additionally comprise one or more other enzymes, such as amylase, lipase, cutinase, another protease, cellulase, 20 peroxidase and oxidase.

The detergent may contain 1-65 % of a detergent builder or complexing agent such as zeolite, diphosphate, triphosphate, phosphonate, citrate, nitrilotriacetic acid (NTA), ethylenediaminetetraacetic acid (EDTA), diethylenetriaminepenta- acetic acid (DTMPA), alkyl- or alkenylsuccinic acid, soluble silicates or layered 25 silicates (e.g. SKS-6 from Hoechst). The detergent may also be unbuilt, i.e. essentially free of detergent builder.

The detergent may comprise one or more polymers. Examples are carboxymethylcellulose (CMC), poly(vinylpyrrolidone) (PVP),

polyethyleneglycol (PEG), poly(vinyl alcohol) (PVA), polycarboxylates such as polyacrylates, maleic/acrylic acid copolymers and lauryl methacrylate/acrylic acid copolymers.

The detergent may contain a bleaching system which may comprise a H₂O₂ source such as perborate or percarbonate which may be combined with a peracid-forming bleach activator such as tetraacetythylenediamine (TAED) or nonanoyloxybenzenesulfonate (NOBS). Alternatively, the bleaching system may comprise peroxyacids of e.g. the amide, imide, or sulfone type.

The enzymes of the detergent composition of the invention may be stabilized using conventional stabilizing agents, e.g. a polyol such as propylene glycol or glycerol, a sugar or sugar alcohol, lactic acid, boric acid, or a boric acid derivative as e.g. an aromatic borate ester, and the composition may be formulated as described in e.g. WO 92/19709 and WO 92/19708.

The detergent may also contain other conventional detergent ingredients such as e.g. fabric conditioners including clays, foam boosters, suds suppressors, anti-corrosion agents, soil-suspending agents, anti-soil redeposition agents, dyes, bactericides, optical brighteners, or perfume.

The pH (measured in aqueous solution at use concentration) will usually be neutral or alkaline, e.g. 7-12 such as 7-10.5.

Particular forms of detergent compositions within the scope of the invention include:

- 1) A detergent composition formulated as a granulate having a bulk density of at least 600 g/l comprising
- linear alkylbenzenesulfonate (calculated as acid) 7 - 12%
 - 5 - alcohol ethoxysulfate (e.g. C₁₂₋₁₈ alcohol, 1-2 EO) or alkyl sulfate (e.g. C₁₈₋₁₈) 1 - 4%
 - alcohol ethoxylate (e.g. C₁₄₋₁₅ alcohol, 7 EO) 5 - 9%
 - 10 - sodium carbonate (as Na₂CO₃) 14 - 20%
 - soluble silicate (as Na₂O, 2SiO₂) 2 - 6%
 - zeolite (as NaAlSiO₄) 15 - 22%
 - sodium sulfate (as Na₂SO₄) 0 - 6%
 - sodium citrate/citric acid 0 - 15%
 - 15 (as C₆H₅Na₃O₇/C₆H₈O₇)
 - sodium perborate (as NaBO₃·H₂O) 11 - 18%
 - TAED 2 - 6%
 - carboxymethylcellulose 0 - 2%
 - polymers (e.g. maleic/acrylic acid copolymer, PVP, PEG) 0 - 3%
 - 20 - enzymes 0 - 5%
 - minor ingredients (e.g. suds suppressors, perfume, optical brightener, photobleach) 0 - 5%
- 25 2) A detergent composition formulated as a granulate having a bulk density of at least 600 g/l comprising
- linear alkylbenzenesulfonate (calculated as acid) 6 - 11%
 - alcohol ethoxysulfate
 - 30 (e.g. C₁₂₋₁₈ alcohol, 1-2 EO)

or alkyl sulfate (e.g. C ₁₆₋₁₈)	1 - 3%
- alcohol ethoxylate (e.g. C ₁₄₋₁₅ alcohol, 7 EO)	5 - 9%
- sodium carbonate (as Na ₂ CO ₃)	15 - 21%
5 - soluble silicate (as Na ₂ O, 2SiO ₂)	1 - 4%
- zeolite (as NaAlSiO ₄)	24 - 34%
- sodium sulfate (as Na ₂ SO ₄)	4 - 10%
- sodium citrate/citric acid (as C ₆ H ₅ Na ₃ O ₇ /C ₆ H ₈ O ₇)	0 - 15 %
10 - carboxymethylcellulose	0 - 2%
- polymers (e.g. maleic/acrylic acid copolymer, PVP, PEG)	1 - 6%
- enzymes	0 - 5%
- minor ingredients	
15 (e.g. suds suppressors, perfume)	0 - 5%

3) A detergent composition formulated as a granulate having a bulk density of at least 600 g/l comprising

- linear alkylbenzenesulfonate (calculated as acid)	5 - 9%
20 - alcohol ethoxylate (e.g. C ₁₂₋₁₅ alcohol, 7 EO)	7 - 14%
- soap as fatty acid (e.g. C ₁₆₋₂₂)	1 - 3%
- sodium carbonate (as Na ₂ CO ₃)	10 - 17%
25 - soluble silicate (as Na ₂ O, 2SiO ₂)	3 - 9%
- zeolite (as NaAlSiO ₄)	23 - 33%
- sodium sulfate (as Na ₂ SO ₄)	0 - 4%
- sodium perborate (as NaBO ₃ ·H ₂ O)	8 - 16%

- TAED	2 - 8%
- phosphonate (e.g. EDTMPA)	0 - 1%
- carboxymethylcellulose	0 - 2%
5 - polymers (e.g. maleic/acrylic acid copolymer, PVP, PEG)	1 - 3%
- enzymes	0 - 5%
- minor ingredients (e.g. suds suppressors, perfume, optical brightener)	0 - 5%

4) A detergent composition formulated as a granulate having a bulk density of
10 at least 600 g/l comprising

- linear alkylbenzenesulfonate (calculated as acid)	8 - 12%
- alcohol ethoxylate (e.g. C ₁₂₋₁₅ alcohol, 7 EO)	10 - 25%
15 - sodium carbonate (as Na ₂ CO ₃)	14 - 22%
- soluble silicate (as Na ₂ O, 2SiO ₂)	1 - 5%
- zeolite (as NaAlSiO ₄)	25 - 35%
- sodium sulfate (as Na ₂ SO ₄)	0 - 10%
- carboxymethylcellulose	0 - 2%
20 - polymers (e.g. maleic/acrylic acid copolymer, PVP, PEG)	1 - 3%
- enzymes	0 - 5%
- minor ingredients (e.g. suds suppressors, perfume)	0 - 5%

25 5) An aqueous liquid detergent composition comprising

- linear alkylbenzenesulfonate

(calculated as acid)	15 - 21%
- alcohol ethoxylate (e.g. C ₁₂₋₁₅ alcohol, 7 EO or C ₁₂₋₁₅ alcohol, 5 EO)	12 - 18%
5 - soap as fatty acid (e.g. oleic acid)	3 - 13%
- alkenylsuccinic acid (C ₁₂₋₁₄)	0 - 13%
- aminoethanol	8 - 18%
- citric acid	2 - 8%
- phosphonate	0 - 3%
10 - polymers (e.g. PVP, PEG)	0 - 3%
- borate (as B ₄ O ₇)	0 - 2%
- ethanol	0 - 3%
- propylene glycol	8 - 14%
- enzymes	0 - 5%
15 - minor ingredients (e.g. dispersants, suds suppressors, perfume, optical brightener)	0 - 5%
6) An aqueous structured liquid detergent composition comprising	
- linear alkylbenzenesulfonate	
20 (calculated as acid)	15 - 21%
- alcohol ethoxylate (e.g. C ₁₂₋₁₅ alcohol, 7 EO or C ₁₂₋₁₅ alcohol, 5 EO)	3 - 9%
- soap as fatty acid (e.g. oleic acid)	3 - 10%
25 - zeolite (as NaAlSiO ₄)	14 - 22%
- potassium citrate	9 - 18%
- borate (as B ₄ O ₇)	0 - 2%

	- carboxymethylcellulose	0 - 2%
	- polymers (e.g. PEG, PVP)	0 - 3%
	- anchoring polymers as e.g. lauryl methacrylate/acrylic acid copolymer;	
5	molar ratio 25:1; MW 3800	0 - 3%
	- glycerol	0 - 5%
	- enzymes	0 - 5%
	- minor ingredients (e.g. dispersants, suds suppressors, perfume,	
10	optical brighteners)	0 - 5%

7) A detergent composition formulated as a granulate having a bulk density of at least 600 g/l comprising

	- fatty alcohol sulfate	5 - 10%
	- ethoxylated fatty acid monoethanolamide	3 - 9%
15	- soap as fatty acid	0 - 3%
	- sodium carbonate (as Na_2CO_3)	5 - 10%
	- soluble silicate (as $\text{Na}_2\text{O}, 2\text{SiO}_2$)	1 - 4%
	- zeolite (as NaAlSiO_4)	20 - 40%
	- sodium sulfate (as Na_2SO_4)	2 - 8%
20	- sodium perborate (as $\text{NaBO}_3 \cdot \text{H}_2\text{O}$)	12 - 18%
	- TAED	2 - 7%
	- polymers (e.g. maleic/acrylic acid copolymer, PEG)	1 - 5%
	- enzymes	0 - 5%
25	- minor ingredients (e.g. optical brightener, suds suppressors, perfume)	0 - 5%

8) A detergent composition formulated as a granulate comprising

- linear alkylbenzenesulfonate (calculated as acid)	8 - 14%
- ethoxylated fatty acid monoethanolamide	5 - 11%
- soap as fatty acid	0 - 3%
5 - sodium carbonate (as Na_2CO_3)	4 - 10%
- soluble silicate (as $\text{Na}_2\text{O}, 2\text{SiO}_2$)	1 - 4%
- zeolite (as NaAlSiO_4)	30 - 50%
- sodium sulfate (as Na_2SO_4)	3 - 11%
- sodium citrate (as $\text{C}_6\text{H}_5\text{Na}_3\text{O}_7$)	5 - 12%
10 - polymers (e.g. PVP, maleic/acrylic acid copolymer, PEG)	1 - 5%
- enzymes	0 - 5%
- minor ingredients (e.g. suds suppressors, perfume)	0 - 5%
15 9) A detergent composition formulated as a granulate comprising	
- linear alkylbenzenesulfonate (calculated as acid)	6 - 12%
- nonionic surfactant,	1 - 4%
- soap as fatty acid	2 - 6%
20 - sodium carbonate (as Na_2CO_3)	14 - 22%
- zeolite (as NaAlSiO_4)	18 - 32%
- sodium sulfate (as Na_2SO_4)	5 - 20%
- sodium citrate (as $\text{C}_6\text{H}_5\text{Na}_3\text{O}_7$)	3 - 8%
- sodium perborate (as $\text{NaBO}_3 \cdot \text{H}_2\text{O}$)	4 - 9%
25 - bleach activator (e.g. NOBS or TAED)	1 - 5%
- carboxymethylcellulose	0 - 2%

- polymers (e.g. polycarboxylate or PEG) 1 - 5%
 - enzymes 0 - 5%
 - minor ingredients
(e.g. optical brightener, perfume) 0 - 5%
- 5' 10) An aqueous liquid detergent composition comprising
- linear alkylbenzenesulfonate
(calculated as acid) 15 - 23%
 - alcohol ethoxysulfate
(e.g. C₁₂₋₁₅ alcohol, 2-3 EO) 8 - 15%
 - 10 - alcohol ethoxylate
(e.g. C₁₂₋₁₅ alcohol, 7 EO
or C₁₂₋₁₅ alcohol, 5 EO) 3 - 9%
 - soap as fatty acid (e.g. lauric acid) 0 - 3%
 - aminoethanol 1 - 5%
 - 15 - sodium citrate 5 - 10%
 - hydrotrope (e.g. sodium toluenesulfonate) 2 - 6%
 - borate (as B₄O₇) 0 - 2%
 - carboxymethylcellulose 0 - 1%
 - ethanol 1 - 3%
 - 20 - propylene glycol 2 - 5%
 - enzymes 0 - 5%
 - minor ingredients (e.g. polymers, dispersants,
perfume, optical brighteners) 0 - 5%
- 11) An aqueous liquid detergent composition comprising
- 25 - linear alkylbenzenesulfonate
(calculated as acid) 20 - 32%
 - alcohol ethoxylate

(e.g. C ₁₂₋₁₅ alcohol, 7 EO or C ₁₂₋₁₅ alcohol, 5 EO)	6 - 12%
- aminoethanol	2 - 6%
- citric acid	8 - 14%
5 - borate (as B ₄ O ₇)	1 - 3%
- polymer (e.g. maleic/acrylic acid copolymer, anchoring polymers as e.g. lauryl methacrylate/acrylic acid copolymer and CMC)	0 - 3%
10 - glycerol	3 - 8%
- enzymes	0 - 5%
- minor ingredients (e.g. hydrotropes, dispersants, perfume, optical brighteners)	0 - 5%
 12) A detergent composition formulated as a granulate having a bulk density 15 of at least 600 g/l comprising	
- anionic surfactant (linear alkylbenzenesulfonate, alkyl sulfate, alpha- olefinsulfonate, alpha-sulfo fatty acid methyl esters, alkanesulfonates, soap)	25 - 40%
20 - nonionic surfactant (e.g. alcohol ethoxylate)	1 - 10%
- sodium carbonate (as Na ₂ CO ₃)	8 - 25%
- soluble silicates (as Na ₂ O, 2SiO ₂)	5 - 15%
- sodium sulfate (as Na ₂ SO ₄)	0 - 5%
25 - zeolite (as NaAlSiO ₄)	15 - 28%
- sodium perborate (as NaBO ₃ ·4H ₂ O)	0 - 20%
- bleach activator (TAED or NOBS)	0 - 5%
- enzymes	0 - 5%

- minor ingredients
(e.g. perfume, optical brighteners) 0 - 3%

13) Detergent formulations as described in 1) - 12) where the content of linear alkylbenzenesulfonate - or a part of it - is substituted by alkyl sulfate (C_{12} - C_{18}).

- 5 14) Detergent formulations as described in 1) - 13) which contain a stabilized or encapsulated peracid either as an additional component or as a substitute for already specified bleach systems.

15) Detergent compositions as described in 3), 7), 9) and 12) where the content of perborate is substituted with percarbonate.

- 10 16) Detergent composition formulated as a nonaqueous detergent liquid comprising a liquid nonionic surfactant as e.g. linear alkoxyated primary alcohol, a builder system (e.g. phosphate), enzyme and alkali. The detergent may also comprise anionic surfactant and/or a bleach system.

- The trypsin-like protease of the invention may be incorporated in
15 concentrations conventionally employed in detergents. It is at present contemplated that, in the detergent composition of the invention, the trypsin-like protease may be added in an amount corresponding to 0.001-100 mg of the enzyme per liter of wash liquor.

BRIEF DESCRIPTION OF THE DRAWINGS

- 20 The invention is further illustrated in the accompanying drawings, in which

Fig. 1 illustrates the construction of the expression plasmid pSX183 further described in Example 3.

The present invention is further illustrated in the following examples which should not, in any manner, be considered to limit the scope of the present invention.

MATERIALS AND METHODS

5 Strains

Aspergillus oryzae IFO 4177

Fusarium oxysporum DSM 2672 described in WO 89/06270

E. coli MC1000 (Cassabadian and Cohen, 1980)

Determination of trypsin-like *Fusarium* protease activity

- 10 The trypsin-like protease is assayed using the specific substrate N-Benzoyl-L-arginine p-nitroanilide hydrochloride (L-BAPA or L-BAPNA).

Buffer

0.01 M dimethylglutaric acid (Sigma D4379), 0.2 M boric acid and 0.002 M calcium chloride adjusted to pH 6.5 with NaOH.

15 Substrate

L-BAPA is available from Sigma (B3133) or Merck (Art.10754).

A 0.2 M stock solution in dimethyl sulfoxide is prepared (87mg/ml) (store frozen) and diluted to 0.004 M with the buffer described above.

Assay

- 20 The trypsin-like protease is diluted to approx. 15 μ g/ml for the assay.

100 μ l of the diluted trypsin-like protease are mixed with 100 μ l of 0.004 M L--BAPA in a 96 well microtiter plate. A blank with 100 μ l buffer and 100 μ l 0.004 M L-BAPA is used for correction.

The absorption change (delta OD/min) is monitored at 405 nm in an Elisa reader for 10 minutes with readings every 5 minutes at 25°C or room temperature.

The result is calculated relative to the trypsin content of a reference *Fusarium* trypsin-like protease.

Plasmids

- 10 pCDV1-PL (Noma et al., 1986)
- p777 (described in EP 0 489 718)
- pToC90 (described in WO 91/17243)

EXAMPLES

EXAMPLE 1

15 PURIFICATION OF THE TRYPSIN-LIKE PROTEASE FROM FUSARIUM

A crude *F. oxysporum* protease preparation was isolated from the supernatant obtained by fermentation of the *F. oxysporum* DSM 2672 in a soy meal containing medium substantially as described in WO 89/06270. Subsequently, the *Fusarium oxysporum* trypsin-like protease was separated from two other
20 *F. oxysporum* proteases by 0-1M/0-25% sodium chloride/isopropanol gradient elution at pH 7.0 from a bacitracin-silicapolyol affinity column (Mortensen et al., 1989). The two other proteases separated from the *F. oxysporum* trypsin-like protease by this procedure were, a subtilisin-like enzyme, and an aspartic protease with specificity required for the activation of the proform of the *F.*
25 *oxysporum* trypsin-like protease (Asn-Ile cleavage).

The other purification steps were carried out in accordance with well-known procedures, and included ion exchange chromatography on Separose CL-6B (prior to the bacitracin separation), gel filtration and concentration by ultrafiltration.

- 5 The specific absorption for a 0.1% solution of the purified trypsin-like *F.oxysporum* protease at 280 nm was determined to 1.32.

The purified protease was subsequently subjected to amino acid sequencing and analysis, and the N-terminal amino acid sequence was determined.

EXAMPLE 2

10 CLONING OF A *FUSARIUM OXYSPORUM* TRYPSIN GENE

mRNA preparation

- Mycelium from a culture of *the F. oxysporum* strain DSM 2672 obtained as described in Example 1 above, was crushed under liquid nitrogen, resuspended in 5M guanidine rhodanide, separated on a CsCl-sarcosyl gradient, and mRNA
15 was isolated therefrom. The mRNA was washed with ethanol, and fractionated on an oligo-dT column (Chirgwin et al. (1978), Truelsen et al. (1979)). The quality of the resulting mRNA was tested in an *in vitro* translation assay using reticulocyte lysate and ³⁵S-methionine. An SDS-PAGE analysis showed bands from below 20 to more than 100 kDa.

20 Preparation and screening of a cDNA library

- A cDNA library of the mRNA was constructed in *E. coli* MC 1000 following the procedure of Noma et al. (1986). After transformation and plating about 15000 colonies were replicated onto Whatman 540 filter paper. A mixed 17-mer probe based on the N-terminal of the enzyme was synthesised (T_M 42-52
25 °C).

26

N-terminal : I V G G T S A G D F P F I V S R N G

Probe : G A T T T T C C N T T T A T C G T

5 T

Split on the isoleucine codons the probe was divided into three pools of 32 and three sets of filters were screened. After wash at 42°C and re-isolation of possible candidates three positive clones were identified. These positive clones were all found with the ATC-isoleucine probe.

10 Identification and analysis of the gene

Plasmids were prepared from the three clones according to Sambrook et al, 1989 and, following restriction mapping, two turned out to be identical and the third was about 100 bp shorter. The longer clones were sequenced on both strands using the method of Maxam & Gilbert (1980) and the resulting

15 DNA sequence appears from SEQ ID No. 1.

The amino acid composition of the purified *F. oxysporum* trypsin-like protease obtained as described in Examples 1 was in accordance with the composition deduced from the DNA-sequence shown in SEQ ID No. 1.

Analysis of this sequence revealed a gene coding for a trypsin-like protease

20 with a 17 amino acid signal peptide according to the rules of von Heijne (1986) (cleaved between Ala-17 and Ala-18) and a propeptide of 7 amino acids (Ala-Pro-Gln-Glu-Ile-Pro-Asn) between the signal peptidase cleavage site and the known N-terminus of the enzyme. The role of this hypothetical propeptide is unknown but proteases are normally made as inactive zymogens.

25 The mature enzyme shows less than 50% identity with all known trypsins be it mammalian, from fruit fly, or bacterial (*Streptomyces*). The highest identity (48%) of the mature trypsin-like protease was found with a fruit fly trypsin.

Expression of the gene product

cDNA encoding the trypsin-like *F. oxysporum* protease, obtained as described above, was inserted into the vector pCDV1-PL described by Noma et al. (1986) resulting in the plasmid pSX180. The coding region of the cDNA was
5 inserted as a NcoI-XbaI fragment into the *Aspergillus* expression plasmid p777 (EP O 489 718) which was cut with BamHI and partially with XbaI. To join the 5'end of the cloned DNA to the vector a synthetic linker DNA KFN709/710 (illustrated in Fig. 1) was added to the ligation reaction. The resulting plasmid pSX183 was co-transformed into *A. oryzae* (IFO 4177) together with plasmid
10 pToC90 carrying the amdS from *A. nidulans* (WO 91/17243). Transformants were selected for growth on acetamide. When grown in YPD medium a protein appeared in the supernatant slightly larger than the trypsin-like protease isolated from *F. oxysporum* supernatants. Surprisingly no activity of the gene product could be determined from hydrolysis of L-Benzoyl-Arginoyl-pNA (a
15 trypsin substrate).

EXAMPLE 3

ACTIVATION OF THE PROENZYME

In an attempt to obtain an active trypsin-like protease, an aspartyl protease isolated from *F. oxysporum* supernatants (as described in Example 1) was
20 added to the fermentation medium harbouring the inactive trypsin-like protease.

By this treatment it was found that the protein from the transformed *A. oryzae* became the same size as the standard on an SDS gel and the sample showed tryptic activity as determined from hydrolysis of L-Benzoyl-Arginoyl-pNA (data
25 not shown). This indicates that the enzyme is expressed in the form of a proenzyme, and thus confirms the presence of a propeptide as deduced from

the DNA sequence. From the activity studies it is apparent that the function of the propeptide is to keep the enzyme in an inactive form.

REFERENCES CITED IN THE SPECIFICATION

- Chirgwin, J. M., Przybyla, A. E., MacDonald, R. J., and Rutter, W. J. (1978). Isolation of biologically active ribonucleic acid from sources enriched in ribonuclease. *Biochemistry* 18: 5294-5299.
- 5 Truelsen, E., Gausing, K., Jochimsen, B., Jørgensen, P., and Marcker, K. A. (1979). Cloning of soybean leghemoglobin structural gene sequences synthesized in vitro. *Nucleic Acids Res.* 6: 3061-3072.
- Christensen, T., Woeldike, H., Boel, E., Mortensen, S. B., Hjortshøj, K., Thim, L., and Hansen, M. T. (1988). High level expression of recombinant genes in Aspergillus
10 oryzae. *Bio/Technology* 6: 1419-1422.
- Maxam, A. M., and Gilbert, W. (1980). Sequencing end-labelled DNA with base-specific chemical cleavages. *Methods Enzymol.* 65: 499-
- von Heijne, G. (1986). A new method for predicting signal sequence cleavage sites. *Nucleic Acids Res.* 14: 4683-4690.
- 15 Hudson, L., and Hay, F. Practical Immunology, Third edition (1989), Blackwell Scientific Publications
- Lipman and Pearson, *Science* 227, 1435 (1985)
- Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor, 1989
- 20 Saiki, R.K. et al., *Science* 239, 1988, pp. 487-491, 1988
- Beaucage et al., *Tetrahedron Letters* 22, 1981, pp. 1859-1869

Matthes et al., The EMBO J. 3, 1984, pp. 801-805

Malardier et al., Gene 78 (1989), pp. 147-156

Zamost et al., Journal of Industrial Microbiology, Vol. 5, pp. 303-312, 1990

Noma et al. (1986), Nature 319, 640-646

5 Cassabadian and Cohen (1980), J.Mol.Biol. 138, 179-203

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

- (A) NAME: NOVO NORDISK A/S
- 5 (B) STREET: Novo Alle
- (C) CITY: Bagsvaerd
- (E) COUNTRY: DENMARK
- (F) POSTAL CODE (ZIP): DK-2880
- (G) TELEPHONE: +45 44448888
- 10 (H) TELEFAX: +45 4449 3256
- (I) TELEX: 37304

(ii) TITLE OF INVENTION: Trypsin-like Fusarium protease

(iii) NUMBER OF SEQUENCES: 2

(iv) COMPUTER READABLE FORM:

- 15 (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (EPO)

20 (2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 998 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- 25 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

30 (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Fusarium oxysporum
- (B) STRAIN: DSM 2672

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- 35 (B) LOCATION: 1..998

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

ATCATCAACC ACTCTTCACT CTTCAACTCT CCTCTCTTGG ATATCTATCT CTTCACCATG	60
GTCAAGTTGG CTTCCGTGGT TGCACTTGTT GCTCCCTTGG CTGCTGCGC TOCTCAGGAG	120
ATCCCCAACA TTGTTGGTGG CACTTCTGOC AGCGCTGGOG ACTTTCCCTT CATGTGAGC	180
5 ATTAGCGCA ACGGTGGGCC CTGGTGTGGA GGTCTCTTCC TCAAOGCAA CACGTCTTG	240
ACTGCTGCC ACTGCGTTTC CGGATACGCT CAGAGCGGT TCCAGATTGG TGCTGGCAGT	300
CTGCTGCGA CTTCTGGTGG TATTACCTCC TCGCTTTCTT CCGTCAGAGT TCACCTAGC	360
TACAGCGGA ACAACAAGA TCTTGCTATT CTGAAGCTCT CTACTTCAT CCGCTCGGC	420
GGAAACATGG GCTATGCTGG CCGGCTGCT TCGGCTCTG ACCGTGTGGC TGGATCTTCT	480
10 GGCACGTGG CTGGCTGGGG CGCTACCTCT GAGGCGGCA GCTCTACTCC CGTCAACCTT	540
CTGAAGGTTA CTGTCCCTAT CGTCTCTGGT GCTACCTGCC GAGCTCAGTA CGGCACCTCC	600
GCCATCAACA ACCAGATGTT CTGTGCTGGT GTTCTTTCCG GTGGCAAGGA CTCTTGCCAG	660
GGTGACAGCG GCGGCCCCAT CGTGACAGC TCCAACACTC TTATCGGTGC TGTCTCTTGG	720
GGTAACGGAT GTGCCCGACC CAACTACTCT GGTGCTATG CCAGCGTTGG TGCTCTCGGC	780
15 TCTTTCATG ACACCTATGC TPAAATACCT TGTTGGAAGC GTGAGATGT TCCTTGAATA	840
TTCTCTAGCT TGAGTCTTGG ATACGAAACC TGTTTGAGAA ATAGGTTTCA ACAGTTAAG	900
AAGATATGAG TTGATTTCAG TTGGATCTTA GTCCGGTGG CTGTAATAG AGCAATCTAG	960
ATAGCCCAAA TTGAATATGA AATTGATGA AAATATTC	998

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 248 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: protein
 (iii) HYPOTHETICAL: NO
 (v) FRAGMENT TYPE: internal
 (vi) ORIGINAL SOURCE:
 (A) ORGANISM: *Fusarium oxysporum*
 (B) STRAIN: DSM 2672

10 (ix) FEATURE:
 (A) NAME/KEY: Protein
 (B) LOCATION: 25..248

(ix) FEATURE:
 (A) NAME/KEY: Peptide
 (B) LOCATION: 1..24
 (D) OTHER INFORMATION: /label= pre-propeptide

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Met Val Lys Phe Ala Ser Val Val Ala Leu Val Ala Pro Leu Ala Ala
 -20 -15 -10

Ala Ala Pro Gln Glu Ile Pro Asn Ile Val Gly Gly Thr Ser Ala Ser
 -5 1 5

Ala Gly Asp Phe Pro Phe Ile Val Ser Ile Ser Arg Asn Gly Gly Pro
 10 15 20

25 Trp Cys Gly Gly Ser Leu Leu Asn Ala Asn Thr Val Leu Thr Ala Ala
 25 30 35 40

His Cys Val Ser Gly Tyr Ala Gln Ser Gly Phe Gln Ile Arg Ala Gly
 45 50 55

Ser Leu Ser Arg Thr Ser Gly Gly Ile Thr Ser Ser Leu Ser Ser Val
 60 65 70

30 Arg Val His Pro Ser Tyr Ser Gly Asn Asn Asn Asp Leu Ala Ile Leu
 75 80 85

Lys Leu Ser Thr Ser Ile Pro Ser Gly Gly Asn Ile Gly Tyr Ala Arg
 90 95 100

Leu Ala Ala Ser Gly Ser Asp Pro Val Ala Gly Ser Ser Ala Thr Val
 105 110 115 120

5 Ala Gly Trp Gly Ala Thr Ser Glu Gly Gly Ser Ser Thr Pro Val Asn
 125 130 135

Leu Leu Lys Val Thr Val Pro Ile Val Ser Arg Ala Thr Cys Arg Ala
 140 145 150

10 Gln Tyr Gly Thr Ser Ala Ile Thr Asn Gln Met Phe Cys Ala Gly Val
 155 160 165

Ser Ser Gly Gly Lys Asp Ser Cys Gln Gly Asp Ser Gly Gly Pro Ile
 170 175 180

Val Asp Ser Ser Asn Thr Leu Ile Gly Ala Val Ser Trp Gly Asn Gly
 185 190 195 200

15 Cys Ala Arg Pro Asn Tyr Ser Gly Val Tyr Ala Ser Val Gly Ala Leu
 205 210 215

Arg Ser Phe Ile Asp Thr Tyr Ala
 220


CLAIMS

1. An active recombinant trypsin-like protease enzyme comprising the amino acid residues 25-224 of the amino acid sequence shown in the appended SEQ ID No. 2.
- 5 2. A proteolytic composition comprising a recombinant protease enzyme as defined in claim 1.
3. A DNA construct comprising a DNA sequence encoding the recombinant protease enzyme as claimed in claim 1.
4. A DNA construct according to claim 3, wherein the DNA sequence is as
10 shown in the appended SEQ ID No. 1.
5. A DNA construct according to any of claims 3 or 4, in which the DNA sequence encoding the recombinant protease enzyme is derived from a microorganism.
6. A DNA construct according to claim 5, in which the DNA sequence is
15 derivable from a bacterium or fungus.
7. A DNA construct according to claim 6, in which the DNA sequence is derivable from a strain of a *Fusarium* species, in particular a strain of *F. oxysporum*.
8. A recombinant expression vector comprising a DNA construct according
20 to any of claims 3-7.
9. A cell comprising a DNA construct according to any of claims 3-7 or a vector according to claim 8.

10. A cell according to claim 9, which is a microbial cell.
11. A cell according to claim 10 which is a bacterial cell or a fungal cell.
12. A cell according to claim 11, in which the bacterial cell is a cell of a gram-positive bacterium, e.g. of the genus *Bacillus* or *Streptomyces* or a cell
5 of a gram-negative bacterium, e.g. of the genus *Escherichia*, and the fungal cell is a yeast cell, e.g. of the genus *Saccharomyces*, or a cell of a filamentous fungus, e.g. of the genus *Aspergillus* or *Fusarium*.
13. A method of producing a recombinant protease enzyme as defined in claim 1, wherein a cell according to any of claims 9-12 is cultured under con-
10 ditions conducive to the production of the protease, and the protease is subsequently recovered from the culture.
14. The method according to claim 13, in which the recombinant protease enzyme is expressed in the form of a proenzyme and the cell is cultured in the presence of a proteolytic enzyme capable of converting the proenzyme of the
15 protease into a mature enzyme.
14. The method according to claim 19, in which the proteolytic enzyme is a metallo-protease, in particular a *Bacillus* metallo-protease.
15. A detergent additive comprising a recombinant protease enzyme according to claim 1, optionally in the form of a non-dusting granulate,
20 stabilised liquid or protected enzyme.
16. A detergent additive according to claim 14 which contains 0.02-200 mg of enzyme protein/g of the additive.

17. A detergent additive according to claim 15 or 16 which additionally comprises another enzyme such as another protease, an amylase, a lipase, a peroxidase and/or a cellulase.
18. A detergent composition comprising a recombinant protease enzyme 5 according to claim 1.
19. A detergent composition according to claim 18 which additionally comprises another enzyme such as another protease, an amylase, a lipase, a peroxidase and/or a cellulase.

1/1



KFN 709/710:

GATCCTCGAATTCTCTTCAGATCTCTTCAC

 GAGCTTAAGAGAAGTCTAGAGAAGTGGTAC

BamHI EcoRI BglII NcoI

FIG. 1

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 94/00177

A. CLASSIFICATION OF SUBJECT MATTER		
IPC 5: C12N 9/76, C12N 15/57, C11D 3/386 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols)		
IPC 5: C12N		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
SE,DK,FI,NO classes as above		
Electronic data base consulted during the international search (names of data base and, where practicable, search terms used)		
BIOSIS, MEDLINE, WPIL, US PATENTS FULLTEXT DATABASES, SEQUENCE DATABASE		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO, A1, 8906270 (NOVO INDUSTRI A/S), 13 July 1989 (13.07.89), page 4, line 8 - line 12; page 4, line 34 - page 5, line 3; page 8, line 4 - line 7, page 8 line 19 - line 21 --	1-19
A	WO, A1, 9300925 (AMGEN INC.), 21 January 1993 (21.01.93), page 7, line 28 - page 8, line 3; page 13, line 14 - line 20; page 16, line 24 - page 17, line 5 --	1-19
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input checked="" type="checkbox"/> See patent family annex.		
* Special categories of cited documents "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
23 August 1994		29 -08- 1994
Name and mailing address of the ISA/ Swedish Patent Office Box 5055, S-102 42 STOCKHOLM Facsimile No. +46 8 666 02 86		Authorized officer Carolina Palmcrantz Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 94/00177

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	PROTEIN ENGINEERING, Volume 6, No 4, 1993, W.R. Rypniewski et al, "The sequence and X-ray structure of the trypsin from Fusarium oxysporum" page 341 - page 348 -- -----	1-19

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 94/00177

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 13
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
The wording "...under conditions conducive to..." does not clearly define which the intended conditions are. Therefore, claim 13 is not considered to be clear and concise according to PCT article 6.
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐

The additional search fees were accompanied by the applicant's protest.

☐

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT
Information on patent family members

02/07/94

International application No.

PCT/DK 94/00177

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A1- 8906270	13/07/89	DE-U- 6890098 EP-A, B- 0394352 EP-A- 0471265	16/04/92 31/10/90 19/02/92
WO-A1- 9300925	21/01/93	AU-A- 2307592 CA-A- 2090463 EP-A- 0546156	11/02/93 02/01/93 16/06/93